



CLIENTS | PEOPLE | PERFORMANCE

Water Corporation

Report for Mundaring Water Treatment Plant Site Selection

Preliminary Spring Flora and Fauna Assessment - **Pine Plantation**

June 2006



Contents

Executive Summary	i
1. Introduction	1
1.1 Background	1
1.2 Scope of this Report	1
1.3 Site Location	2
1.4 Limitations of this Survey	2
2. Desktop Assessment	4
2.1 Vegetation	4
2.2 Vegetation Extent and Status	4
2.3 Threatened Ecological Communities	5
2.4 Flora	6
2.5 Fauna	8
3. Field Survey	14
3.1 Survey Methods	14
3.2 Vegetation Types	14
3.3 Vegetation Condition	14
3.4 Threatened Ecological Communities	15
3.5 Flora	16
3.6 Fauna	16
4. Conclusions and Recommendations	18
5. Limitations	20
6. References	21

Table Index

Table 1	Vegetation extent and status for Beard (1979) Vegetation Associations in the Mundaring area (after Shepherd, <i>et al.</i> (2002), Shepherd, pers. comm.)	5
Table 2	Significant Flora Species Previously Recorded within the Mundaring Area (Source: DEC and the WA Herbarium)	6



Table 3	Listing of Potentially Occurring Rare and Priority Fauna Species – Mundaring Weir area (Source: <i>EPBC Act</i> Protected Matters Search, DEC Threatened and Priority Fauna Database search results and WA Museum Faunabase search for the general Mundaring Weir area)	9
Table 4	Government of Western Australia (2000) Vegetation Condition Scale	15
Table 5	Conservation Codes and Descriptions for DEC Declared Rare and Priority Flora Species.	25
Table 6	Flora List for the ‘Pine Plantation’ Site	26
Table 7	<i>Environment Protection And Biodiversity Conservation Act 1999</i> Fauna Conservation Categories	31
Table 8	<i>Western Australian Wildlife Conservation Act 1950</i> Fauna Conservation Codes	32
Table 9	DEC Priority Fauna Conservation Codes. (Species not listed under the Wildlife Conservation Act 1950, but for which there is some concern).	33
Table 10	Fauna Species that may occur in the Pine Plantation Site (Source: Western Australian Museum <i>Faunabase</i> Search of the Mundaring area); and those recorded during the opportunistic fauna surveys.	34

Figure Index

Figure 1	Location Plan
Figure 2	Vegetation Types at the Pine Plantation Site
Figure 3	Vegetation Condition at the Pine Plantation Site

Appendices

- A Figures
- B Flora
- C Vegetation
- D Fauna



Executive Summary

The Water Corporation is currently conducting a site selection process for the proposed Mundaring Water Treatment Plant (WTP). This process involves community-based consultation, and has included preliminary consideration of the environmental, social, technical and economic features of a number of site options.

As part of the site selection process GHD Pty Ltd (GHD) was commissioned by the Water Corporation to undertake preliminary flora and fauna investigations in Spring 2006 at some of the Mundaring WTP site options. This report presents the results of the field survey at the Pine Plantation Site.

This flora and fauna investigation is a preliminary investigation aimed at identifying any major environmental constraints on the site and to assess the environmental values of the site. A full flora and vegetation assessment and fauna assessment will be required once a final site has been determined.

The preliminary flora and fauna investigations for the Pine Plantation Site included a desktop assessment and a preliminary field assessment. The results of the assessment concluded that:

- » The site is within State Forest and contains a pine plantation.
- » This site has been previously logged and subject to other disturbances. Approximately half of the site contains regrowth Jarrah-Marri forest but the other half has been converted to pine plantation. The Jarrah-Marri forest occurs in the north-west part of the site and the pine plantation in the south-east of the site. There is also Jarrah-Marri forest adjacent to the access track (approximately 1km long) and there will be clearing of this native vegetation required for the upgrade of the access track and laying of pipes into the site.
- » The Jarrah-Marri forest at this site is mainly Condition 2/3 (*Excellent/Very Good*). However, there are some patches of vegetation that are more highly disturbed, including an area that has been used as a dump which is Condition 6 (*Completely degraded*). The pine plantation was generally Condition 5 (*Degraded*).
- » The Site has moderate species diversity with a total of 74 taxa (59 native taxa) from 27 families recorded within the surveyed area. The majority of the species occurred within the area of native vegetation in the north-west of the site.
- » No Declared Rare or Priority Species were identified during the survey.
- » An eastern states grass species, *Austrodanthonia fulva*, was recorded in the survey which has not been recorded in WA before. This species was probably introduced from planes using the nearby landing strip.
- » A total of 15 weed species were recorded within the Site, which represents about 20 % of the total species recorded. These were mainly pasture grasses (Poaceae) and *Acacia* species (Mimosaceae) from the eastern states. Around half of the site (the pine plantation) contains predominantly introduced species



- » A number of fauna species occur in the general Mundaring area, including a number of protected species. Of significance, noted in the opportunistic survey, were Carnaby's Cockatoos, a species listed as Endangered under the *Wildlife Conservation Act 1950* and the *Environment Protection and Biodiversity Conservation Act*.
- » The habitat value of the site was assessed during the survey. This site contains areas of native vegetation that may be used by native fauna species; however, this habitat is fairly uniform and has been disturbed in the past and cannot be considered of 'high value'. The pine plantation may also be used by fauna species but is of less value than that of the surrounding native vegetation.
- » There is an existing road to this site, which could be upgraded and used as the access route for the Mundaring WTP. The upgrading of this approximately 1km of track will include clearing of native vegetation and laying of pipelines. The access track and pipelines has the potential to impact on fauna linkages.

Approximately half of the site contains regrowth native vegetation and the other half contains pine plantation. The impacts on flora and fauna of a WTP at this site are not “significant” and could be managed. However, clearing of native vegetation will be required, particularly along the access track.



1. Introduction

1.1 Background

The Water Corporation proposes to improve the quality and security of water supplies to the Eastern Hills areas of Perth and to the Goldfields and Agriculture regions. The \$55 million Mundaring Water Supply Improvement Project was announced in 2002, and involved the first stage construction over about six years of a 50 megalitre water tank and a 150 megalitres per day capacity water treatment plant at Sawyers Valley Tank site. Two more stages are planned and by 2030 it is projected that a total of three 50 megalitre water tanks would be constructed on the site and the water treatment plant capacity would be increased to 250 megalitres per day.

Strong community objections emerged against the location of the Water Treatment Plant (WTP) at Sawyers Valley. Due to these community objections the Water Corporation decided to withdraw the application to the EPA to build the water treatment plant at Sawyers Valley and a commitment was made to undergo a more detailed site selection process, with the involvement of the community. The Water Corporation's application to the EPA to build one 50 megalitre water tank remained active.

The current site selection process commenced in mid-2005 and is a community-based consultation process, which involved reconsidering all previous site options and including any other potentially suitable options. This process has involved preliminary consideration of the environmental, social, technical and economic features of a number of site options.

1.2 Scope of this Report

GHD Pty Ltd (GHD) has been commissioned by the Water Corporation to assist with the site selection process for the Mundaring WTP and to provide independent advice to the Water Corporation and the community on the environmental features of the site options, as required.

As part of the site selection process GHD was commissioned by the Water Corporation to undertake preliminary biological field surveys in Spring 2006 at a number of the Mundaring WTP site options. A number of site options being considered at this time contained major areas of remnant vegetation. These included:

- » Site One;
- » Site Two;
- » Site '1km below Weir;
- » Pine Plantation Site;
- » DEC Land (formerly 'CALM Land);
- » Site Five (Sawyers Hill); and
- » Site Seven.



There have been previous biological investigations conducted at Site One and Two, so further survey was not considered necessary at this time. A number of biological surveys have been conducted previously at Site Five but none were conducted in early Spring so a survey of this site was considered necessary at this time. A biological survey was not undertaken at Site Seven but a brief inspection of the site was conducted to determine the environmental condition of the site.

The Water Corporation determined that preliminary biological surveys in spring 2006 were to be conducted at the following sites:

- » Site '1km below Weir';
- » Pine Plantation Site;
- » DEC Land (formerly 'CALM Site'); and
- » Site Five (Sawyers Hill).

This report presents the results of the field survey at the Site 'Pine Plantation', (**Figure 1**).

The biological investigations were designed to highlight major flora and fauna issues and to determine the ecological value of the site. The surveys including the following aspects:

- » A broad-scale vegetation assessment, including assessment of vegetation types, vegetation condition, dominant species and potential presence of diseases such as dieback, and weeds. The vegetation type and condition was mapped;
- » A targeted search for any Declared Rare and Priority Flora that occur in the area;
- » An assessment of habitat potential, including the assessment of the significance of habitat trees; and
- » A targeted opportunistic fauna search, particularly for any significant fauna species.

1.3 Site Location

The Site 'Pine Plantation' is a 20 ha site located in State forest on Allen Rd to the north of Mundaring Weir, off Mundaring Weir Road. This site is a pine plantation next to the DEC Airstrip north of Mundaring Weir and also includes some adjacent native vegetation.

This site is very close to Bibbulmun Track, the track passes through the north-west corner of site.

1.4 Limitations of this Survey

This biological investigation is only a preliminary survey and a full flora and vegetation assessment and fauna assessment may be required once a final site has been determined. The investigation focused on the site of the WTP and detailed investigations of the access routes was not undertaken. However, the potential access routes were assessed to determine any major biological constraints.



The targeted Declared Rare and Priority Flora search is dependent on a number of factors, including flowering times for the significant species, and hence cannot be a completely comprehensive search. Further field-work may be required once the final site has been selected to determine the potential presence of any significant flora species at that site.

Desktop investigations on fauna species were conducted at the regional level, to determine species that are present in the general area and that have the potential to occur at the site. While this does not allow comparison between sites the information on fauna habitat obtained during the survey provides some indication of the likelihood of fauna species occurring at the site and of the value of the site as fauna habitat.



2. Desktop Assessment

2.1 Vegetation

Beard (1979) has mapped the vegetation of the general area at a scale of 1:250,000. The site vegetation is within the Darling Botanical District: Dale Subdistrict of the South Western Botanical Province as recognised by Beard (1979). According to Beard the vegetation of the sub-district is primarily Jarrah – Marri forest.

Mapping by Heddle *et al.* (1980) shows the site to be within the **Dwellingup Complex in Medium to high rainfall**. Open forest of *Eucalyptus marginata* – *E. calophylla* (now *Corymbia calophylla*).

Mundaring Weir and the study site occur on the Darling Plateau region, which has the following general vegetation complexes as defined in the Regional Forest Agreement Vegetation Complexes mapping of Matiske and Havel (1998):

- » **Dwellingup 2 (D2) (uplands)** Open forest of *Eucalyptus marginata subsp. marginata* – *Corymbia calophylla* on lateritic uplands in the arid zone.
- » **Helena 2 (He2) (in valleys)** Mosaic of open forest of *Eucalyptus marginata subsp. thalassica*-*Corymbia calophylla* and woodland of *Eucalyptus wandoo* with some *Eucalyptus accedens* and *Eucalyptus rudis* on the deeper soils ranging to closed heaths and lithic complex on shallow soils associated with granite on steep slopes of valleys in semiarid and arid zones.
- » **Murray 2 (My2) (in valleys)** Open forest of *Eucalyptus marginata subsp. thalassica*-*Corymbia calophylla*- *Eucalyptus patens* and woodlands of *Eucalyptus wandoo* with some *Eucalyptus accedens* on valley slopes to woodlands of *Eucalyptus rudis*-*Melaleuca raphiophylla* on the valley floor in semiarid and arid zones.
- » **Yarragil 1 (in valleys)** Open forest of *Eucalyptus marginata subsp. marginata*-*Corymbia calophylla* on slopes with mixtures of *Eucalyptus patens* and *Eucalyptus megacarpa* on the valley floors in humid and subhumid zones.
- » **Pindalup (in valleys)** Open forest of *Eucalyptus marginata subsp. thalassica*-*Corymbia calophylla* on slopes and open woodlands of *Eucalyptus wandoo* with some *Eucalyptus patens* on the lower slopes in semiarid and arid zones.

The Matiske and Havel mapping (1998) indicates that the 'Pine Plantation' site occurs mainly in the vegetation complex **Dwellingup 2 (D2) (uplands)**.

2.2 Vegetation Extent and Status

A vegetation type is considered underrepresented if there is less than 30 percent of its original distribution remaining. From a purely biodiversity perspective, and not taking into account any other land degradation issues, there are several key criteria now being applied to vegetation in States where clearing is still occurring (EPA, 2000)

- » The “threshold level” below which species loss appears to accelerate exponentially at an ecosystem level is regarded as being at a level of 30% of the pre-European/ pre-1750 extent of the vegetation type;
- » A level of 10% of the original extent is regarded as being a level representing *Endangered*; and
- » Clearing which would put the threat level into the class below should be avoided.

Such status can be delineated into five (5) classes, where:

- » *Presumed Extinct*: Probably no longer present in the bioregion
- » *Endangered**: <10% of pre-European extent remains
- » *Vulnerable**: 10-30% of pre-European extent exists
- » *Depleted**: >30% and up to 50% of pre-European extent exists
- » *Least Concern*: >50% pre-European extent exists and subject to little or no degradation over a majority of this area.

* or a combination of depletion, loss of quality, current threats and rarity gives a comparable status

Native vegetation types represented in the survey area, their regional extent and reservation status are generally drawn from Shepherd, *et al.* (2002), and Shepherd pers. comm., which are in turn based on broadscale mapping undertaken by Beard (1979). These are shown in **Table 1**.

Table 1 Vegetation extent and status for Beard (1979) Vegetation Associations in the Mundaring area (after Shepherd, *et al.* (2002), Shepherd, pers. comm.)

Vegetation Association	Vegetation Description	Pre-European Extent (ha)	Current Extent (ha)	% Remaining
3	Medium Forest: Jarrah – Marri	908, 040	747, 888	82.4

Generally the extent of the vegetation remaining on the Darling Scarp is above the threshold level and it can be seen from the table above the Jarrah – Marri Medium Forest is relatively well preserved.

2.3 Threatened Ecological Communities

Ecological communities are defined as ‘naturally occurring biological assemblages that occur in a particular type of habitat’ (English and Blythe, 1997). Threatened Ecological Communities (TECs) are ecological communities that have been assessed and assigned to one of four categories related to the status of the threat to the community, i.e. Presumed Totally Destroyed, Critically Endangered, Endangered, and Vulnerable. Some TECs are protected under the Commonwealth *Environment Protection and Biodiversity Act 1999* (EPBC Act). Although TECs are not formally protected under the State *Wildlife Conservation Act 1950*, the loss of, or disturbance to, some TECs triggers the EPBC Act. The Environmental Protection Authority’s (EPA’s) position on



TECs states that proposals that result in the direct loss of TECs are likely to require formal assessment.

A search of the DEC's TEC database was undertaken for the Mundaring area. This search recorded that there were no known occurrences of TECs for the general area of Mundaring Weir. Occurrences of the 'Critically Endangered' communities known as 'Shrublands and woodlands of the eastern side of the Swan Coastal Plain' and '*Eucalyptus calophylla* – *Xanthorrhoea preissii* woodlands and shrublands, Swan Coastal Plain' and an occurrence of the 'Endangered' threatened ecological community known as '*Banksia attenuata* woodland over species rich dense shrublands' occur nearby to the general Mundaring area, and the potential for these communities to occur in the study site was taken into consideration during the field surveys.

2.4 Flora

Flora species that are considered to be significant are listed under the Western Australian *Wildlife Conservation Act 1950* and the Commonwealth *EPBC Act*. A description of the DEC's conservation codes is provided in **Table 5, Appendix B**.

A search of the DEC's Rare Flora Databases and the Western Australian Herbarium indicates that a number of significant species are known to occur in the general Mundaring area. There are no records of significant species at or near to the site, but this may simply reflect the lack of flora surveys conducted in the area. The recorded significant species in the general vicinity of the site are presented in **Table 2** and **Figure 1, Appendix A**. Please note that a large search area was requested due to the large study area for all the Mundaring WTP site options, and this large search area has resulted in a large number of recorded significant flora species.

Table 2 Significant Flora Species Previously Recorded within the Mundaring Area (Source: DEC and the WA Herbarium)

Species	Conservation Code
<i>Adenanthos cygnorum</i> subsp. <i>chamaephyton</i>	P3
<i>Adenanthos filifolius</i>	P3
<i>Acacia anomala</i>	DRF
<i>Acacia aphylla</i>	DRF
<i>Acacia horridula</i>	P3
<i>Acacia oncinophylla</i> subsp. <i>oncinophylla</i>	P3
<i>Anthocercis gracilis</i>	DRF
<i>Aotus cordifolia</i>	P3
<i>Boronia tenuis</i>	P4
<i>Bossiaea modesta</i>	P2



<i>Caladenia arrecta</i>	P4
<i>Cyanicula ixiooides</i> subsp. <i>ixiooides</i>	P4
<i>Darwinia apiculata</i>	DRF
<i>Darwinia pimelioides</i>	P4
<i>Daviesia oxylobium</i>	P4
<i>Diplolaena andrewsii</i>	P2
<i>Grevillea drummondii</i>	
<i>Grevillea flexuosa</i>	DRF
<i>Grevillea pimeleoides</i>	P4
<i>Halgania corymbosa</i>	P3
<i>Hemigenia rigida</i>	P1
<i>Isopogon drummondii</i>	P3
<i>Lasiopetalum bracteatum</i>	P4
<i>Lepyrodia heleocharoides</i>	P3
<i>Persoonia sulcata</i>	P4
<i>Pimelea rara</i>	P4
<i>Pithocarpa corymbulosa</i>	P2
<i>Scholtzia</i> sp. Bickley	P1
<i>Senecio gilbertii</i>	P1
<i>Senecio leucoglossus</i>	P4
<i>Stylidium</i> sp. Boulder Rock	
<i>Stylidium striatum</i>	P4
<i>Templetonia drummondii</i>	P4
<i>Tetralthea pilifera</i>	P3
<i>Tetralthea</i> sp. Granite	P3
<i>Thelymitra manginiorum</i>	DRF
<i>Thysanotus anceps</i>	P3
<i>Verticordia lindleyi</i> subsp. <i>lindleyi</i>	P4
<i>Verticordia serrata</i> var. <i>linearis</i>	P3



2.5 Fauna

A search of the Western Australian Museum records was undertaken, which determined that 19 amphibian species, 46 bird species, 26 mammal species, 3 fish species and 56 reptile species have been recorded within the general Mundaring area. These results are included in **Table 10, Appendix D**. Please note that a large area was searched for fauna records, due to the large study area for all the Mundaring WTP site options, and this large search area has resulted in a substantial number of recorded fauna species.

2.5.1 Fauna of Conservation Significance

The conservation status of fauna species is assessed under State and Commonwealth Acts; in particular the Western Australian *Wildlife Conservation Act 1950*; *Wildlife Conservation (Specially Protected Fauna) Notice 2006*, and the Commonwealth *EPBC Act*.

The significance levels for fauna used in the *EPBC Act* are those recommended by the International Union for the Conservation of Nature and Natural Resources (IUCN). These levels are described in **Table 7, Appendix D**.

The *EPBC Act* also lists migratory species that are recognised under international treaties such as the China Australia Migratory Bird Agreement (CAMBA), the Japan Australia Migratory Bird Agreement (JAMBA) and the Bonn Convention (The Convention on the Conservation of Migratory Species of Wild Animals).

Listed migratory species also include any native species identified in an international agreement approved by the Commonwealth Environment Minister. The Minister may approve an international agreement for this purpose if satisfied that it is an agreement relevant to the conservation of migratory species.

In Western Australia, the *Wildlife Conservation (Specially Protected Fauna) Notice 2006* has significant levels for fauna classified in a series of Schedules (**Table 8, Appendix C**). The DEC also produces a supplementary list of Priority Fauna, being species that are not considered Threatened under the Western Australian *Wildlife Conservation Act 1950* but for which the Department feels there is a cause for concern. These species have no special protection, but their presence would normally be considered and they need further survey and evaluation of conservation status before consideration can be given to declaration as threatened fauna. Levels of Priority are described in **Table 9, Appendix C**.

A listing of Significant Fauna from the *EPBC Act* Protected Matters Search Tool and Rare and Priority species from the Department of Environment and Conservation (DEC) rare fauna database for the general Mundaring area are presented in **Table 3**.

It should be noted that the results of the searches are only an indication of the threatened fauna that may be present in the general area and detailed fauna surveys, including trapping programs are required to verify these results.



Table 3 Listing of Potentially Occurring Rare and Priority Fauna Species – Mundaring Weir area (Source: EPBC Act Protected Matters Search, DEC Threatened and Priority Fauna Database search results and WA Museum Faunabase search for the general Mundaring Weir area)

Genus	Species	Common Name	DEC/ Wildlife Conservation Act 1950 Rating	EPBC Act Rating	CALM Database	EPBC Act Protected Matters search
<i>Dasyurus</i>	<i>geoffroi</i>	Chuditch	Vulnerable / Schedule 1	Vulnerable	x	
<i>Myrmecobius</i>	<i>fasciatus</i>	Numbat	Vulnerable / Schedule 1	Vulnerable	x	
<i>Calyptorhynchus</i>	<i>latirostris</i>	Carnaby's Cockatoo	Endangered / Schedule 1	Endangered	x	
<i>Calyptorhynchus</i>	<i>baudinii</i>	Baudin's Cockatoo	Endangered / Schedule 1	Vulnerable		x
<i>Pseudocheirus</i>	<i>occidentalis</i>	Western Ringtail Possum	Vulnerable / Schedule 1	Vulnerable		X
<i>Phascogale</i>	<i>tapoatafa tapoatafa</i>	Southern Brush-tailed Phascogale	Priority 3		x	
<i>Macropus</i>	<i>irma</i>	Western Brush Wallaby	Priority 4		x	
<i>Hydromys</i>	<i>chrysogaster</i>	Water-rat (Rakali)	Priority 4		x	
<i>Isoodon</i>	<i>obesulus fusciventer</i>	Quenda	Priority 5		x	
<i>Haliaeetus</i>	<i>leucogaster</i>	White-bellied Sea-eagle		Migratory		X
<i>Apus</i>	<i>pacificus</i>	Fork-tailed Swift		Marine		X



Genus	Species	Common Name	DEC/ Wildlife Conservation Act 1950 Rating	EPBC Act Rating	CALM Database	EPBC Act Protected Matters search
<i>Ardea</i>	<i>alba</i>	Great Egret, White Egret		Marine		X
<i>Ardea</i>	<i>ibis</i>	Cattle Egret		Marine		X
<i>Merops</i>	<i>ornatus</i>	Rainbow Bee-eater		Marine		X



Some species that appear in the *EPBC Act* Protected Matters Search Tool are often not likely to occur within the specified area, as the search provides an approximate guidance to matters of national significance that require further investigation. The records from the DEC searches of threatened fauna provide more accurate information for the general area, however some records of sightings or trappings can be dated and often misrepresent the current range of threatened species.

The habitat requirements of the protected species listed under the *EPBC Act*, the *Wildlife Conservation Act 1950* and listed as priority species by the DEC, and the likelihood of their occurrence in the site is considered below.

***Dasyurus geoffroii* Chuditch**

The Chuditch is the largest carnivorous marsupial in Western Australia. This species occupies a wide range of habitats including woodlands, riparian vegetation, beaches and deserts. The Chuditch formerly ranged over nearly 70 % of Australia but now retains only a patchy distribution through the Jarrah forest and mixed Karri/Marri/Jarrah forest of south-western WA. This reduction in range and decline in population numbers have been caused by habitat alteration, impacts from the introduction of foxes and cats, hunting and poisoning. This species is currently listed as Vulnerable on the *EPBC Act* (CALM, 2005a).

The site contains some habitat that is suitable for Chuditch and they have been known to occur in the region, so there is the potential for this species to utilise the site.

***Myrmecobius fasciatus* Numbat**

The Numbat is a small, banded, diurnal marsupial that feeds solely on termites. This species once ranged widely in southern semi-arid and arid Australia, distributed within a number of vegetation types. However, the Numbat's current distribution is limited to Dryandra and Perup/Kingston area east of Manjimup and a number of nature reserves into which it has been reintroduced. This species occupies a number of habitat types including Jarrah forest, open eucalypt woodland, banksia woodland and tall closed shrubland (CALM, 2005b).

This species only occurs in isolated populations in Western Australia, in a number of small reserves, and it is unlikely that this species would be found at this site.

***Calyptorhynchus latirostris* Carnaby's Cockatoo**

Carnaby's Cockatoo, also known as the Short-billed Black-Cockatoo, is distributed across the south-west of Western Australia in uncleared or remnant areas of eucalypt woodland and shrubland or kwongan heath. Breeding usually occurs in the Wheatbelt region of Western Australia, with flocks moving to the higher rainfall coastal areas to forage after the breeding season. These black cockatoos feed on the seeds of a variety of native plants, including *Allocasuarina*, *Banksia*, *Dryandra*, *Eucalyptus*, *Grevillea* and *Hakea*, and some introduced plants, including *Pinus*. They will also feed on the nectar from flowers of a number of species, and on insect larvae. Carnaby's Cockatoo has been listed as Endangered under the Federal *EPBC Act*.



This species is known to occur in the study area and could use a number of the plant species on this site for foraging.

***Calyptrorhynchus baudinii* Baudin's Cockatoo**

Baudin's Cockatoo, also known as the Long-billed Black-Cockatoo, is found in the south-west of Western Australia in the forest and woodlands of Jarrah (*Eucalyptus marginata*), Karri (*E. diversicolor*) and Marri (*Corymbia calophylla*). The primary food source of this cockatoo is the seeds of the Marri (Garnett and Crowley, 2000). This species has been impacted by the removal of large Marri throughout its range. Baudin's Cockatoo has been listed as Vulnerable under the Commonwealth *EPBC Act*.

Baudin's Cockatoo are known to occur in the Mundaring area and the site contains habitat that is suitable for this species. This species may utilise the site for foraging, and the site also contains Marri trees, that are used by this cockatoo for breeding.

***Pseudocheirus occidentalis* Western Ringtail Possum**

The Western Ringtail Possum (*Pseudocheirus occidentalis*) is found in the south-west of Western Australia, with a patchy occurrence from the Collie River to Two Peoples Bay (Jones, *et al.*, 1994). Urban populations of this species are located in Busselton, as well as East Augusta and Albany.

This species was recorded in the *EPBC Act* Protected Matters Search but is not known from the Mundaring area and this record is likely to be inaccurate.

***Phascogale tapoatafa tapoatafa* Southern Brush-tailed Phascogale**

This arboreal marsupial occurs in forest and woodland with suitable tree hollows. Populations fluctuate dramatically in response to invertebrate prey abundance. This subspecies has been observed in dry sclerophyll forests and open woodlands that contain hollow-bearing trees but a sparse ground cover. Records are less common from wetter forests (CALM, 2005c).

The Southern Brush-tailed Phascogale is known to occur in low densities in the northern Jarrah forests. This species is known to occur in the Mundaring area and may occur on the study site.

***Macropus irma* Western Brush Wallaby**

The Western Brush Wallaby, a medium sized macropod, is a grazer found primarily in open forest and woodland. This species was once very common in the south-west of Western Australia but has undergone a reduction in range and a significant decline in abundance in its current habitat. The decline in populations of this species has resulted from extensive clearing within its original distribution and from predation of juvenile Western Brush Wallabies by foxes (CALM, 2005d).

This species is known to occur in the general area and may occur on the study site.

***Hydromys chrysogaster* Water-rat (Rakali)**

The water rat occupies habitat in the vicinity of permanent water and nests are constructed in logs or at the end of tunnels dug into banks. Unlike many other Australian rodents, the water rat is not entirely nocturnal, with activity usually high at



sunset, though animals have been seen foraging during the day. The water rat is an opportunistic predator, feeding upon large aquatic insects, fish, crustaceans and mussels. They are also known to feed on frogs, lizards, small mammals, fresh carrion, and birds (CALM, 2005e).

This species may occur in the general Mundaring area but there is no suitable habitat for this species at the study site and it would be unlikely to occur here.

***Isodon obesulus fusciventer* Quenda**

The Quenda is an omnivorous marsupial that occurs in the south-west of Western Australia. It prefers areas with dense understorey vegetation, particularly around swamps and along watercourses. However, it also occurs in woodlands, and may use less ideal habitat where this habitat occurs adjacent to the thicker, more desirable vegetation. On the Swan Coastal Plain Quenda are often associated with wetlands (CALM, 2005c).

Quenda are known to occur in the Mundaring region and may be present at the site.



3. Field Survey

3.1 Survey Methods

The field survey was conducted on the 27th September, 2006, by a qualified botanist and a field ecologist.

The survey recorded vegetation types and dominant species in each vegetation type. A full list of species was generated for the site and where identification was uncertain, confirmation was made at the Western Australian State Herbarium. The presence of DRF and Priority flora was assessed. The condition and weed status of the vegetation were also noted. Vegetation was also assessed to determine the presence of TECs within the project area. Aerial photography was used to assist in the delineation of vegetation types present in the study area.

A reconnaissance survey was conducted for the presence of fauna. A consideration of fauna habitat was undertaken within the study area.

Nomenclature of the species follows that of *Florabase* (2006) for flora species and the *Faunabase* (2006) for fauna species.

3.2 Vegetation Types

The vegetation at this site is fairly uniform, with two main vegetation types occurring; a Jarrah-Marri forest in the north-west part of the site and a Pine plantation in the south-east of the site. A full description of the vegetation types, photographs and inventory of vascular plants is provided in **Appendix B** and **Appendix C**. The broad vegetation types at the site have been mapped in **Figure 2, Appendix A**.

The Jarrah-Marri forest in the north-west of the site is fairly uniform, but shows some variation between the upland areas (in the west of this vegetation) and the flatter sections in the east. The upland areas contain scattered Bull Banksia (*Banksia grandis*) and Sheoak (*Allocasuarina fraseriana*), which are mainly lacking in the lower sections. There are patches of vegetation in this area that have evidence of disturbance, including scattered Pine trees, and a dominance of Parrot Bush (*Dryandra sessilis*).

The pine plantation is dominated by Pine trees but also contains introduced Golden Wattle (*Acacia pycnantha*) and scattered young *Eucalyptus* species.

3.3 Vegetation Condition

The vegetation in the study area was given a condition rating based on the Bush Forever (Government of Western Australia, 2000) vegetation condition ratings scale. This scale recognises a level of intactness of vegetation, which is defined by the following:

- » Completeness of structural levels;
- » Extent of weed invasion;



- » Historical disturbance from tracks and other clearing or dumping;
- » The potential for natural or assisted regeneration.

The ratings in this scale are described as follows:

Table 4 Government of Western Australia (2000) Vegetation Condition Scale

Assigned Number	Classification	Description
1	<i>Pristine or nearly so</i>	No obvious signs of disturbance
2	<i>Excellent</i>	Vegetation structure intact, disturbance affecting individual species, and weeds are non-aggressive species
3	<i>Very Good</i>	Vegetation structure altered, obvious signs of disturbance
4	<i>Good</i>	Vegetation structure significantly altered by very obvious signs of multiple disturbance, retains basic vegetation structure or ability to regenerate it
5	<i>Degraded</i>	Basic vegetation structure severely impacted by disturbance. Scope for regeneration but not to a state approaching good condition without intensive management.
6	<i>Completely Degraded</i>	The structure of the vegetation is no longer intact and the area is completely or almost without native species

The Jarrah-Marri forest at this site is mainly Condition 2/3 (*Excellent/Very Good*). However, there are some patches of vegetation that are more highly disturbed, including an area that has been used as a dump which was Condition 6 (*Completely degraded*). The Pine plantation was generally Condition 5 (*Degraded*) as it contained predominantly introduced species, with some scattered natives. The vegetation condition of the site is recorded at **Figure 3**.

3.3.1 Plant Pests and Diseases

There was some evidence to suggest that *Phytophthora cinnamomi* (commonly known as dieback) was present within the Site, based on patterns of death of susceptible plants. Should this site be considered further for the Mundaring WTP the determination of the presence of dieback should be assessed formally by a DEC interpreter or equivalent.

3.4 Threatened Ecological Communities

No Threatened Ecological Communities were identified during the field survey.



3.5 Flora

The Site has moderate species diversity with a total of 74 taxa (59 native taxa) from 27 families recorded within the surveyed area.

The dominant families recorded from the area were:

- » Proteaceae 9 taxa
- » Myrtaceae 7 taxa
- » Mimosaceae 6 taxa

Additionally, the dominant genus recorded was:

- » *Acacia* 6 taxa
- » *Stylidium* 4 taxa

Eight species were identified to genus only, due to lack of distinctive features, such as flowers. Additionally one species was identifiable to family only, as the majority of the plant had died. Field surveys at different times of year would allow the identification of a greater number of plant species. A number of plants were not identified as they were known to be weeds or garden escapes and were not significant for the survey.

See **Appendix B** for a full list of flora species recorded in the project area.

3.5.1 Significant Species

No Declared Rare or Priority Species were identified during the survey.

3.5.2 Weeds

Weed species were mostly evident along the road that passes through the site, in the pine plantation and at an old dump site on the edge of the road. There was also some weed species scattered throughout the area of native vegetation.

A total of 15 weed species were recorded within the Site, which represents about 20 % of the total species recorded. These were mainly pasture grasses (Poaceae) and *Acacia* species (Mimosaceae) from the eastern states. An eastern states grass species, *Austrodanthonia fulva*, was recorded that has not been recorded in WA before. This species was probably introduced from planes using the adjacent landing strip.

3.6 Fauna

3.6.1 Fauna Species

An opportunistic fauna survey was completed in association with the Spring flora survey. The results from this survey are included in **Table 10, Appendix D**.

Of significance, noted in the opportunistic survey were Carnaby's Cockatoos, a species listed as Endangered under the *Wildlife Conservation Act 1950* and the *EPBC Act*. There was evidence that this species uses the site for foraging.



The number of species determined during the opportunistic species was limited by the short period of the survey and the fact that it was purely opportunistic and did not provide the opportunity to survey those species that are cryptic or not active during the day. Further detailed fauna surveys will be required once the final site of the Mundaring WTP has been selected.

3.6.2 Habitat Value

The habitat value of the site was assessed during the survey. This site contains areas of native vegetation that may be used by native fauna species; however, this habitat is fairly uniform and has been disturbed in the past and cannot be considered of 'high value'. The pine plantation may also be used by fauna species but is of less value than that of the surrounding native vegetation.

3.6.3 Habitat Linkages

Fauna corridors and habitat linkage are important to allow animals to move between areas of resource availability. Such corridors are important for ground and aerial fauna, providing cover, resources, and linking areas suitable for rest and reproduction.

This site would be used for fauna movement; however, as there is good quality bushland on all sides of the site the disruption to this fauna movement would not be significant as the species could still pass around the site.

This site will require upgrading of around 1km of access track into the site, which will include clearing of native vegetation and laying of pipelines. This access track and pipelines has the potential to impact on fauna linkages as some species may be reluctant to cross this track.



4. Conclusions and Recommendations

The preliminary flora and fauna investigations for the Pine Plantation Site included a desktop assessment and a preliminary field assessment. The results of the assessment concluded that:

- » The site is within State Forest and contains a pine plantation.
- » This site has been previously logged and subject to other disturbances. Approximately half of the site contains regrowth Jarrah-Marri forest but the other half has been converted to pine plantation. The Jarrah-Marri forest occurs in the north-west part of the site and the pine plantation in the south-east of the site. There is also Jarrah-Marri forest adjacent to the access track (approximately 1km long) and there will be clearing of this native vegetation required for the upgrade of the access track and laying of pipes into the site.
- » The Jarrah-Marri forest at this site is mainly Condition 2/3 (*Excellent/Very Good*). However, there are some patches of vegetation that are more highly disturbed, including an area that has been used as a dump which is Condition 6 (*Completely degraded*). The pine plantation was generally Condition 5 (*Degraded*).
- » The Site has moderate species diversity with a total of 74 taxa (59 native taxa) from 27 families recorded within the surveyed area. The majority of the species occurred within the area of native vegetation in the north-west of the site.
- » No DRF or Priority flora species were identified during the survey.
- » An eastern states grass species, *Austrodanthonia fulva*, was recorded in the survey that has not been recorded in WA before. This species was probably introduced from planes using the nearby landing strip.
- » A total of 15 weed species were recorded within the Site, which represents about 20 % of the total species recorded. These were mainly pasture grasses (Poaceae) and *Acacia* species (Mimosaceae) from the eastern states. Around half of the site (the pine plantation) contains predominantly introduced species
- » A number of fauna species occur in the general Mundaring area, including a number of protected species. Of significance, noted in the opportunistic survey, were Carnaby's Cockatoos, a species listed as Endangered under the *Wildlife Conservation Act 1950* and the *EPBC Act*.
- » The habitat value of the site was assessed during the survey. This site contains areas of native vegetation that may be used by native fauna species; however, this habitat is fairly uniform and has been disturbed in the past and cannot be considered of 'high value'. The pine plantation may also be used by fauna species but is of less value than that of the surrounding native vegetation.
- » There is an existing road to this site, which could be upgraded and used as the access route for the Mundaring WTP. The upgrading of this approximately 1km of track will include clearing of native vegetation and laying of pipelines. The access track and pipelines has the potential to impact on fauna linkages.



Approximately half of the site contains regrowth native vegetation and the other half contains pine plantation. The impacts on flora and fauna of a WTP at this site are not “significant” and could be managed. However, clearing of native vegetation will be required, particularly along the access track.



5. Limitations

This report presents the results of a Preliminary Flora and Fauna Assessment prepared for the purpose of this commission. The data and advice provided herein relate only to the project described herein and must be reviewed by a competent scientist before being used for any other purpose. GHD accepts no responsibility for other use of the data.

Where reports, searches, any third party information and similar work have been performed and recorded by others the data is included and used in the form provided by others. The responsibility for the accuracy of such data remains with the issuing authority, not with GHD.

For these investigations GHD has conducted desktop data searches and field surveys. The conclusions of this report were based on the information gathered during these investigations and thus reflect the environment of the site at the time of survey. GHD accepts no responsibility for any variation in the flora and fauna present at the site due to natural and seasonal variability.



6. References

- Beard, J.S. (1979). *Vegetation Survey of Western Australia, 1:250,000 Series*. The Vegetation of the Perth Area, Western Australia. Vegmap Publications, Applecross.
- Department of Conservation and Land Management (CALM) (2005a) *Fauna Species Profiles: Chuditch (Dasyurus geoffroii)* [Internet] Available from: http://www.naturebase.net/plants_animals/pdf_files/sp_chuditch.pdf [Accessed 23/11/05]
- Department of Conservation and Land Management (CALM) (2005b) *Fauna Species Profiles: Numbat (Myrmecobius fasciatus)* [Internet] Available from: www.calm.wa.gov.au/plants_animals/pdf_files/sp_numbat.pdf [Accessed 23/11/05]
- Department of Conservation and Land Management (CALM) (2005c) *Fauna Species Profiles: Brush-tailed Phascogale (Phascogale tapoatafa)* [Internet] Available from: http://www.calm.wa.gov.au/plants_animals/pdf_files/sp_brush-tailed_phascogale.pdf [Accessed 23/11/05]
- Department of Conservation and Land Management (CALM) (2005d) *Fauna Species Profiles: Western Brush Wallaby (Macropus irma)*. [Internet] Available from: www.calm.wa.gov.au/plants_animals/pdf_files/sp_western_brush_wallaby.pdf [Accessed 14/10/05]
- Department of Conservation and Land Management (CALM) (2005e). *Fauna Species Profiles: Water Rat (Rakali) (Hydromys chrysogaster)*. Web page access, May 25, 2005. http://www.calm.wa.gov.au/plants_animals/pdf_files/sp_water_rat.pdf
- Department of Conservation and Land Management (CALM) (2005f) *Fauna Species Profiles: Quenda (Isodoon obesulus)*. [Internet] Available from: www.calm.wa.gov.au/plants_animals/pdf_files/sp_quenda.pdf [Accessed 14/10/05]
- Department of Environment and Conservation (DEC) (2006) *Florabase* [Internet] Available from <http://www.calm.wa.gov.au/florabase/index.html>
- English, V and Blythe, J. (1997). *Identifying and Conserving Threatened Ecological Communities in the South West Botanical Province*. Unpublished report for the Department of Conservation and Land Management to Environment Australia.
- Environmental Protection Authority (EPA) (2000) *Protection of Native Vegetation in Western Australia: Clearing of native vegetation, with particular reference to the agricultural area. Position Statement No.2*. December 2000.
- Garnett, S.T. and Crowley, G.M. (2000) *The Action Plan for Australian Birds 2000*. Environment Australia, Canberra, ACT.
- Government of WA, (2000). *Bush Forever Volume 1. Policies, Principles, Processes*. Department of Environmental Protection, Perth, Western Australia.



- Hedde, E.M., Loneragan, O.W. and Havel, J.J. (1980). Vegetation Complexes of the Darling System, Western Australia. In: Atlas of Natural Resources, Darling System Western Australia. Department of Conservation and Environment. WA.
- Jones, B.A., How, R.A. and Kitchener, D.J. (1994b). A Field Study of *Pseudocheirus occidentalis* (Marsupialia:Petauridae). II. Population Studies, *Wildlife Research*: 21, pp 189-201.
- Mattiske, E.M., and Havel, J.J. (1998). *Vegetation Mapping in the South West of Western Australia*. Department of Conservation and Land Management, Perth.
- Orell, P. and Morris, K. (1992) Chuditch Recovery Plan 1992-2001 [Internet] Available from:
<http://www.deh.gov.au/biodiversity/threatened/publications/recovery/chuditch/>
[Accessed 14/11/05]
- Western Australian Museum (2006) *Faunabase* [Internet] Available from
<http://www.museum.wa.gov.au/faunabase/prod/index.htm>



Appendix A

Figures

Figure 1 **Location Plan**

Figure 2 **Vegetation Types at the Pine Plantation Site**

Figure 3 **Vegetation Condition at the Pine Plantation Site**



Appendix B

Flora

Conservation Codes and Descriptions for DEC Declared Rare
and Priority Flora Species.

Flora List for the Pine Plantation Site



Table 5 Conservation Codes and Descriptions for DEC Declared Rare and Priority Flora Species.

Conservation Code	Description
R: Declared Rare Flora – Extant Taxa	Taxa which have been adequately searched for and are deemed to be in the wild either rare, in danger of extinction, or otherwise in need of special protection, and have been gazetted as such.
P1: Priority One – Poorly Known Taxa	Taxa which are known from one or a few (generally <5) populations which are under threat, either due to small population size, or being on lands under immediate threat, e.g. road verges, urban areas, farmland, active mineral leases, etc., or the plants are under threat, e.g. from disease, grazing by feral animals etc. May include taxa with threatened populations on protected lands. Such taxa are under consideration for declaration as 'rare flora', but are in urgent need of further survey.
P2: Priority Two – Poorly Known Taxa	Taxa which are known from one or a few (generally <5) populations, at least some of which are not believed to be under immediate threat (i.e. not currently endangered). Such taxa are under consideration for declaration as 'rare flora', but are in urgent need of further survey.
P3: Priority Three – Poorly Known Taxa	Taxa which are known from several populations, and the taxa are not believed to be under immediate threat (i.e. not currently endangered), either due to the number of known populations (generally >5), or known populations being large, and either widespread or protected. Such taxa are under consideration for declaration as 'rare flora' but are in need of further survey.
P4: Priority Four – Taxa in need of monitoring	Taxa which are considered to have been adequately surveyed and which, whilst being rare (in Australia), are not currently threatened by any identifiable factors. These taxa require monitoring every 5 – 10 years.



Table 6 Flora List for the 'Pine Plantation' Site

Family	Genus	Species	Common Name	Status
Apiaceae	<i>Pentapeltis</i>	<i>peltigera</i>		
Apiaceae	<i>Platysace</i>	<i>filiformis</i>		
Apiaceae	<i>Xanthosia</i>	<i>candida</i>		
Asteraceae	<i>Lagenophora</i>	<i>huegelii</i>		
Asteraceae	<i>Rhodanthe</i>	<i>citrina</i>		
Asteraceae	<i>Senecio</i>	<i>hispidulus</i>	Hispid Fireweed	
Asteraceae	<i>Senecio</i>	<i>pinnatifolius</i>		
Casuarinaceae	<i>Allocasuarina</i>	<i>fraseriana</i>	Sheoak	
Crassulaceae	<i>Crassula</i>	<i>colorata</i>	Dense Stonecrop	
Crassulaceae	<i>Crassula</i>	<i>decumbens</i>	Rufous Stonecrop	
Cyperaceae	<i>Isolepis</i>	<i>marginata</i>	Coarse Club-rush	
Dilleniaceae	<i>Hibbertia</i>	<i>amplexicaulis</i>		
Dilleniaceae	<i>Hibbertia</i>	<i>commutata</i>		
Dilleniaceae	<i>Hibbertia</i>	<i>hypericoides</i>	Yellow Buttercups	
Droseraceae	<i>Drosera</i>	<i>erythrorhiza</i>	Red Ink Sundew	
Droseraceae	<i>Drosera</i>	<i>pallida</i>	Pale Rainbow	
Epacridaceae	<i>Astroloma</i>	<i>ciliatum</i>	Candle Cranberry	
Epacridaceae	<i>Astroloma</i>	<i>sp.</i>		
Epacridaceae	<i>Leucopogon</i>	<i>verticillatus</i>	Tassel Flower	
Euphorbiaceae	<i>Phyllanthus</i>	<i>calycinus</i>	False Boronia	
Goodeniaceae	<i>Lechenaultia</i>	<i>biloba</i>	Blue Leschenaultia	
Iridaceae	<i>Orthrosanthus</i>	<i>laxus</i>	Morning Iris	
Mimosaceae	<i>Acacia</i>	<i>baileyana</i>		*
Mimosaceae	<i>Acacia</i>	<i>dealbata</i>		*
Mimosaceae	<i>Acacia</i>	<i>extensa</i>	Wiry Wattle	
Mimosaceae	<i>Acacia</i>	<i>pycnantha</i>	Golden Wattle	*
Mimosaceae	<i>Acacia</i>	<i>podalyriifolia</i>		*
Mimosaceae	<i>Acacia</i>	<i>saligna</i>		
Myrtaceae	<i>Astartea</i>	<i>sp.</i>		
Myrtaceae	<i>Corymbia</i>	<i>calophylla</i>	Marri	
Myrtaceae	<i>Eucalyptus</i>	<i>marginata</i>	Jarrah	



Family	Genus	Species	Common Name	Status
Myrtaceae	<i>Hypocalymma</i>	<i>angustifolium</i>	White Myrtle	
Myrtaceae	<i>Kunzea</i>	<i>micrantha</i>		
Myrtaceae	<i>Pericalymma</i>	<i>ellipticum</i>	Swamp Teatree	
Myrtaceae				
Orchidaceae	<i>Caladenia</i>	<i>flava</i>	Cowslip Orchid	
Orchidaceae	<i>Disa</i>	<i>bracteata</i>		*
Orchidaceae	<i>Pterostylis</i>	<i>?barbata</i>	?Bird Orchid	
Orchidaceae	<i>Pterostylis</i>	<i>sp.</i>		
Orchidaceae	<i>Thelymitra</i>	<i>sp.</i>		
Oxalidaceae	<i>Oxalis</i>	<i>glabra</i>		*
Oxalidaceae	<i>Oxalis</i>	<i>pes-caprae</i>	Soursob	*
Papilionaceae	<i>Bossiaea</i>	<i>pulchella</i>		
Papilionaceae	<i>Daviesia</i>	<i>decurrens</i>	Prickly Bitter-pea	
Papilionaceae	<i>Kennedia</i>	<i>prostrata</i>	Scarlet Runner	
Papilionaceae	<i>Lupin</i>	<i>sp.</i>		*
Papilionaceae	<i>Trifolium</i>	<i>campestre</i>	Hop Clover	*
Pinaceae	<i>Pinus</i>	<i>radiata</i>		*
Poaceae	<i>Austrodanthonia</i>	<i>fulva</i>		*new record, probably introduced
Poaceae	<i>Briza</i>	<i>minor</i>	Shivery Grass	*
Poaceae	<i>Ehrharta</i>	<i>erecta</i>	Panic Veldt Grass	*
Primulaceae	<i>Anagallis</i>	<i>arvensis</i>	Pimpernel	*
Proteaceae	<i>Adenanthos</i>	<i>barbiger</i>		
Proteaceae	<i>Banksia</i>	<i>grandis</i>	Bull Banksia	
Proteaceae	<i>Dryandra</i>	<i>lindleyana</i>	Couch Honeypot	
Proteaceae	<i>Dryandra</i>	<i>sessilis</i>	Parrot Bush	
Proteaceae	<i>Dryandra</i>	<i>?bipinnatifida</i>		
Proteaceae	<i>Hakea</i>	<i>undulata</i>	Wavy-leaved Hakea	
Proteaceae	<i>Hakea</i>	<i>amplexicaulis</i>	Prickly Hakea	
Proteaceae	<i>Persoonia</i>	<i>elliptica</i>	Spreading Snottygobble	
Proteaceae	<i>Synaphea</i>	<i>acutiloba</i>	Granite Synaphea	
Ranunculaceae	<i>Clematis</i>	<i>sp.</i>		



Family	Genus	Species	Common Name	Status
Ranunculaceae	<i>Clematis</i>	<i>aristata</i>	Clematis	
Rubiaceae	<i>Opercularia</i>	<i>hispidula</i>	Hispid Stinkweed	
Scrophulariaceae	<i>Kickxia</i>	<i>spuria</i>	Roundleaf Toadflax	*
Stylidiaceae	<i>Stylidium</i>	<i>amoenum</i>	Lovely Triggerplant	
Stylidiaceae	<i>Stylidium</i>	<i>calcaratum</i>	Book Triggerplant	
Stylidiaceae	<i>Stylidium</i>	<i>sp.</i>		
Stylidiaceae	<i>Stylidium</i>	<i>hispidum</i>	White Butterfly Triggerplant	
Thymelaeaceae	<i>Pimelea</i>	<i>ciliata</i> subsp. <i>ciliata</i>		
Thymelaeaceae	<i>Pimelea</i>	<i>sp.</i>		
Xanthorrhoea	<i>Xanthorrhoea</i>	<i>gracilis</i>	Graceful Grass Tree	
Xanthorrhoea	<i>Xanthorrhoea</i>	<i>preissii</i>	Grasstree	
Zamiaceae	<i>Macrozamia</i>	<i>riedlei</i>	Zamia	

* Introduced species

NF No flowers (so identification to species was not possible)



Appendix C
Vegetation

Vegetation Community Descriptions at the Site



Appendix D

Fauna

*Environment Protection And Biodiversity Conservation Act
1999 Fauna Conservation Categories*

*Western Australian Wildlife Conservation Act 1950 Fauna
Conservation Codes*

DEC Priority Fauna Conservation Codes.

Fauna Species that may occur in the Pine Plantation Site



Table 7 Environment Protection And Biodiversity Conservation Act 1999 Fauna Conservation Categories

Conservation Category	Definition
Extinct	Taxa not definitely located in the wild during the past 50 years
Extinct in the Wild	Taxa known to survive only in captivity
Critically Endangered	Taxa facing an extremely high risk of extinction in the wild in the immediate future
Endangered	Taxa facing a very high risk of extinction in the wild in the near future
Vulnerable	Taxa facing a high risk of extinction in the wild in the medium-term
Near Threatened	Taxa that risk becoming Vulnerable in the wild
Conservation Dependent	Taxa whose survival depends upon ongoing conservation measures. Without these measures, a conservation dependent taxon would be classified as Vulnerable or more severely threatened.
Data Deficient (Insufficiently Know)	Taxa suspected of being Rare, Vulnerable or Endangered, but whose true status cannot be determined without more information.
Least Concern	Taxa that are not considered Threatened

EPBC Act Categories

An action has, will have, or is likely to have a significant impact on a critically endangered or endangered species if it does, will, or is likely to:

- » Lead to a long-term decrease in the size of a population, or
- » Reduce the area of occupancy of the species, or
- » Fragment an existing population into two or more populations, or
- » Adversely affect habitat critical to the survival of a species, or
- » Disrupt the breeding cycle of a population, or
- » Modify, destroy, remove, isolate or decrease the availability or quality of habitat to the extent that the species is likely to decline, or
- » Result in invasive species that are harmful to a critically endangered or endangered species becoming established in the endangered or critically endangered species' habitat*, or
- » Interfere with the recovery of the species.



** Introducing an invasive species into the habitat may result in that species becoming established. An invasive species may harm a critically endangered or endangered species by direct competition, modification of habitat, or predation.*

An action has, will have, or is likely to have a significant impact on a vulnerable species if it does, will, or is likely to:

- » Lead to a long-term decrease in the size of an important population of a species, or
- » Reduce the area of occupancy of an important population, or
- » Fragment an existing important population into two or more populations, or
- » Adversely affect habitat critical to the survival of a species, or
- » Disrupt the breeding cycle of an important population, or
- » Modify, destroy, remove or isolate or decrease the availability or quality of habitat to the extent that the species is likely to decline, or
- » Result in invasive species that are harmful a vulnerable species becoming established in the vulnerable species' habitat*, or
- » Interferes substantially with the recovery of the species.

An important population is one that is necessary for a species' long-term survival and recovery. This may include populations that are:

- » Key source populations either for breeding or dispersal,
- » Populations that are necessary for maintaining genetic diversity, and/or
- » Populations that are near the limit of the species range.

** Introducing an invasive species into the habitat may result in that species becoming established. An invasive species may harm a vulnerable species by direct competition, modification of habitat, or predation.*

Table 8 Western Australian Wildlife Conservation Act 1950 Fauna Conservation Codes

Conservation Code	Description
Schedule 1	"...fauna that is rare or likely to become extinct, are declared to be fauna that is in need of special protection."
Schedule 2	"...fauna that is presumed to be extinct, are declared to be fauna that is in need of special protection."
Schedule 3	"...birds that are subject to an agreement between the governments of Australia and Japan relating to the protection of migratory birds and birds in danger of extinction, are declared to be fauna that is in need of special protection."
Schedule 4	"...fauna that is in need of special protection, otherwise than for the reasons mentioned [in Schedule 1 – 3]"



Table 9 DEC Priority Fauna Conservation Codes. (Species not listed under the Wildlife Conservation Act 1950, but for which there is some concern).

Conservation Code	Description
Priority 1	Taxa with few, poorly known populations on threatened lands.
Priority 2	Taxa with few, poorly known populations on conservation lands. Taxa which are known from few specimens or sight records from one or a few localities on lands not under immediate threat of habitat destruction or degradation, e.g. national parks, conservation parks, nature reserves, State forest, vacant Crown Land, water reserves, etc.
Priority 3	Taxa which are known from few specimens or sight records, some of which are on lands not under immediate threat of habitat destruction or degradation.
Priority 4	Rare taxa. Taxa which are considered to have been adequately surveyed and which, whilst being rare (in Australia), are not currently threatened by any identifiable factors. These taxa require monitoring every 5 – 10 years.
Priority 5	Taxa in need of monitoring. Taxa which are not considered threatened but are subject to a specific conservation program, the cessation of which would result in the species becoming threatened within five years.



Table 10 Fauna Species that may occur in the Pine Plantation Site (Source: Western Australian Museum *Faunabase* Search of the Mundaring area); and those recorded during the opportunistic fauna surveys.

Family	Genus	Species	Common Name	Status	Recorded During the Field Survey
Birds					
Accipitridae	<i>Accipiter</i>	<i>fasciatus fasciatus</i>	Brown Goshawk		
Accipitridae	<i>Aquila</i>	<i>audax</i>	Wedge-tailed Eagle		
Aegothelidae	<i>Aegotheles</i>	<i>cristatus cristatus</i>	Australian Owlet Nightjar		
Anatidae	<i>Chenonetta</i>	<i>jubata</i>	Australian Wood Duck		
Ardeidae	<i>Ixobrychus</i>	<i>minutus dubius</i>	Little Bittern	P2	
Campephagidae	<i>Coracina</i>	<i>maxima</i>	Ground Cuckoo-shrike		
Corvidae	<i>Corvus</i>	<i>bennetti</i>	Little Crow		
Cracticidae	<i>Cracticus</i>	<i>tibicen dorsalis</i>	Australian Magpie		+
Cracticidae	<i>Strepera</i>	<i>versicolor</i>	Grey Currawong		
Cuculidae	<i>Chrysococcyx</i>	<i>lucidus plagosus</i>	Shining Bronze-cuckoo		
Cuculidae	<i>Cuculus</i>	<i>pallidus</i>	Pallid Cuckoo		
Dicruridae	<i>Rhipidura</i>	<i>fuliginosa preissi</i>	Grey Fantail		
Falconidae	<i>Falco</i>	<i>berigora berigora</i>	Brown Falcon		
Falconidae	<i>Falco</i>	<i>longipennis longipennis</i>	Australian Hobby		
Halcyonidae	<i>Dacelo</i>	<i>novaeguineae</i>	Laughing Kookaburra	*	
Hirundinidae	<i>Hirundo</i>	<i>nigricans</i>	Tree Martin		
Maluridae	<i>Malurus</i>	<i>elegans</i>	Red-winged Fairy-wren		
Meliphagidae	<i>Acanthorhynchus</i>	<i>supercilius</i>	Western Spinebill		
Meliphagidae	<i>Lichmera</i>	<i>indistincta</i>	Brown Honeyeater		+



Family	Genus	Species	Common Name	Status	Recorded During the Field Survey
Motacillidae	<i>Anthus</i>	<i>australis australis</i>	Richard's Pipit		
Pachycephalidae	<i>Pachycephala</i>	<i>pectoralis fuliginosa</i>	Golden Whistler		+
Pardalotidae	<i>Pardalotus</i>	<i>punctatus punctatus</i>	Spotted Pardalote		
Pardalotidae	<i>Pardalotus</i>	<i>punctatus xanthopyge</i>	Yellow-rumped Pardalote		
Pardalotidae	<i>Pardalotus</i>	<i>striatus</i>	Striated Pardalote		
Passeridae	<i>Neochmia</i>	<i>temporalis temporalis</i>	Red-browed Finch		
Passeridae	<i>Stagonopleura</i>	<i>oculata</i>	Red-eared Firetail		
Petroicidae	<i>Petroica</i>	<i>multicolor campbelli</i>	Scarlet Robin		
Phasianidae	<i>Pavo</i>	<i>cristatus</i>	Common Peafowl (Indian Peafowl)	*	
Podargidae	<i>Podargus</i>	<i>strigoides brachypterus</i>	Tawny Frogmouth		
Psittacidae	<i>Calyptorhynchus</i>	<i>banksii</i>	Red-tailed Black Cockatoo		
Psittacidae	<i>Calyptorhynchus</i>	<i>banksii naso</i>	Forest Red-tailed Black Cockatoo	Schedule 1 / Vulnerable (CALM)	
Psittacidae	<i>Calyptorhynchus</i>	<i>baudinii</i>	Baudin's Cockatoo	Schedule 1 / Endangered (CALM) (EPBC - Vulnerable)	
Psittacidae	<i>Calyptorhynchus</i>	<i>latirostris</i>	Carnaby's Cockatoo	Schedule 1 / Endangered	+
Psittacidae	<i>Calyptorhynchus</i>	<i>sp</i>			
Psittacidae	<i>Melopsittacus</i>	<i>undulatus</i>	Budgerigar		
Psittacidae	<i>Neophema</i>	<i>elegans</i>	Elegant Parrot		
Psittacidae	<i>Platycercus</i>	<i>icterotis</i>	Western Rosella		
Psittacidae	<i>Platycercus</i>	<i>spurius</i>	Red-capped Parrot		?



Family	Genus	Species	Common Name	Status	Recorded During the Field Survey
Psittacidae	<i>Platycercus</i>	<i>zonarius</i>	Australian Ringneck (Ring-necked Parrot)		+
Psittacidae	<i>Polytelis</i>	<i>anthopeplus anthopeplus</i>	Regent Parrot		
Rallidae	<i>Porzana</i>	<i>pusilla palustris</i>	Baillon's Crake		
Rallidae	<i>Porzana</i>	<i>tabuensis</i>	Spotless Crake		
Strigidae	<i>Ninox</i>	<i>connivens</i>	Barking Owl	P2 (<i>N. connivens connivens</i>)	
Strigidae	<i>Ninox</i>	<i>novaeseelandiae</i>	Boobook Owl		
Strigidae	<i>Ninox</i>	<i>novaeseelandiae boobook</i>	Boobook Owl		
Turnicidae	<i>Turnix</i>	<i>varia varia</i>	Painted Bustard-Quail		
Mammals					
Burramyidae	<i>Cercartetus</i>	<i>concinnus</i>	Western Pygmy-possum / Mundarda		
Dasyuridae	<i>Antechinus</i>	<i>flavipes</i>	Yellow-footed Antechinus		
Dasyuridae	<i>Antechinus</i>	<i>flavipes leucogaster</i>	Mardo		
Dasyuridae	<i>Dasyurus</i>	<i>geoffroii</i>	Western Quoll / Chuditch	Schedule 1 / Vulnerable	
Dasyuridae	<i>Phascogale</i>	<i>tapoatafa tapoatafa</i>	Brush-tailed Phascogale / Wambenger	P3	
Dasyuridae	<i>Sminthopsis</i>	<i>gilberti</i>	Gilbert's Dunnart		
Felidae	<i>Felis</i>	<i>catus</i>	Cat	*	
Macropodidae	<i>Macropus</i>	<i>eugenii derbianus</i>	Tammar Wallaby	P5	
Macropodidae	<i>Macropus</i>	<i>fuliginosus</i>	Western Grey Kangaroo		+
Macropodidae	<i>Macropus</i>	<i>irma</i>	Western Brush Wallaby	P4	
Macropodidae	<i>Setonix</i>	<i>brachyurus</i>	Quokka	Schedule 1 / Vulnerable	



Family	Genus	Species	Common Name	Status	Recorded During the Field Survey
Muridae	<i>Hydromys</i>	<i>chrysogaster</i>	Water Rat	P4	
Muridae	<i>Mus</i>	<i>musculus</i>	House Mouse	*	
Muridae	<i>Rattus</i>	<i>rattus</i>	Black Rat	*	
Mustelidae	<i>Mustela</i>	<i>putorius</i>	European Polecat / Ferret	*	
Myrmecobiidae	<i>Myrmecobius</i>	<i>fasciatus</i>	Numbat / Walpurti	Schedule 1 / Vulnerable	
Peramelidae	<i>Isoodon</i>	<i>obesulus fusciventer</i>	Quenda / Southern Brown Bandicoot	P5	
Phalangeridae	<i>Trichosurus</i>	<i>vulpecula vulpecula</i>	Common Brushtail Possum		
Tachyglossidae	<i>Tachyglossus</i>	<i>aculeatus</i>	Short-beaked Echidna		
Tarsipedidae	<i>Tarsipes</i>	<i>rostratus</i>	Honey Possum / Noolbenger		
Thylacomyidae	<i>Macrotis</i>	<i>lagotis</i>	Bilby	Schedule 1 / Vulnerable	
Vespertilionidae	<i>Chalinolobus</i>	<i>gouldii</i>	Gould's Wattled Bat		
Vespertilionidae	<i>Nyctophilus</i>	<i>geoffroyi</i>	Lesser Long-eared Bat		
Vespertilionidae	<i>Nyctophilus</i>	<i>timoriensis timoriensis</i>	Greater Long-eared Bat		
Vespertilionidae	<i>Vespadelus</i>	<i>regulus</i>	Southern Forest Bat		
Reptiles					
Agamidae	<i>Ctenophorus</i>	<i>ornatus</i>	Ornate Rock Dragon		
Agamidae	<i>Pogona</i>	<i>minor minor</i>	Western Bearded Dragon		
Agamidae	<i>Rankinia</i>	<i>adelaidensis</i>	Western Heath Dragon		
Boidae	<i>Antaresia</i>	<i>stimsoni stimsoni</i>	Western Stimson's Python		
Boidae	<i>Morelia</i>	<i>spilota imbricata</i>	Southern Carpet Python	Schedule 4	



Family	Genus	Species	Common Name	Status	Recorded During the Field Survey
Cheluidae	<i>Chelodina</i>	<i>oblonga</i>	Oblong Turtle		
Elapidae	<i>Acanthophis</i>	<i>antarcticus</i>	Southern Death Adder	P3	
Elapidae	<i>Brachyuropsis</i>	<i>semifasciata</i>	Southern Shovel-nosed Snake		
Elapidae	<i>Notechis</i>	<i>scutatus</i>	Tiger Snake		
Elapidae	<i>Parasuta</i>	<i>gouldii</i>	Gould's Snake		
Elapidae	<i>Parasuta</i>	<i>nigriceps</i>	Black-backed Snake		
Elapidae	<i>Pseudechis</i>	<i>australis</i>	Mulga Snake		
Elapidae	<i>Pseudonaja</i>	<i>affinis affinis</i>	Dugite		
Elapidae	<i>Pseudonaja</i>	<i>modesta</i>	Ringed Brown Snake		
Elapidae	<i>Pseudonaja</i>	<i>nuchalis</i>	Gwardar		
Gekkonidae	<i>Christinus</i>	<i>marmoratus</i>	Marbled Gecko		
Gekkonidae	<i>Diplodactylus</i>	<i>granariensis granariensis</i>	Wheatbelt Stone Gecko		
Gekkonidae	<i>Diplodactylus</i>	<i>polyopthalmus</i>	Speckled Stone Gecko		
Gekkonidae	<i>Gehyra</i>	<i>variegata</i>	Variegated Tree Dtella		
Gekkonidae	<i>Strophurus</i>	<i>spinigerus</i>	Western Spiny-tailed Gecko		
Gekkonidae	<i>Strophurus</i>	<i>spinigerus inornatus</i>	Western Spiny-tailed Gecko		
Gekkonidae	<i>Underwoodisaurus</i>	<i>millii</i>	Barking Gecko		
Pygopodidae	<i>Aprasia</i>	<i>pulchella</i>	Granite Worm Lizard		
Pygopodidae	<i>Aprasia</i>	<i>repens</i>	Sandplain Worm Lizard		
Pygopodidae	<i>Lialis</i>	<i>burtonis</i>	Burton's Legless Lizard		
Pygopodidae	<i>Pygopus</i>	<i>lepidopodus</i>	Common Scaly-foot		



Family	Genus	Species	Common Name	Status	Recorded During the Field Survey
Scincidae	<i>Acritoscincus</i>	<i>trilineatum</i>	Southwestern Cool Skink		
Scincidae	<i>Cryptoblepharus</i>	<i>plagiocephalus</i>	Fence or Wall Skink		
Scincidae	<i>Ctenotus</i>	<i>delli</i>	Dell's Skink	P4	
Scincidae	<i>Ctenotus</i>	<i>fallens</i>	-		
Scincidae	<i>Ctenotus</i>	<i>labillardieri</i>	Red-legged Skink		
Scincidae	<i>Egernia</i>	<i>kingii</i>	King's Skink		
Scincidae	<i>Egernia</i>	<i>napoleonis</i>	Southwestern Crevice Skink		
Scincidae	<i>Eremiascincus</i>	<i>richardsonii</i>	Banded Skink		
Scincidae	<i>Hemiergis</i>	<i>initialis</i>	Five-toed Earless Skink		
Scincidae	<i>Hemiergis</i>	<i>initialis initialis</i>	Five-toed Earless Skink		
Scincidae	<i>Lerista</i>	<i>distinguenda</i>	-		
Scincidae	<i>Lerista</i>	<i>elegans</i>	-		
Scincidae	<i>Lerista</i>	<i>microtis</i>	-		
Scincidae	<i>Menetia</i>	<i>greyii</i>	Common Dwarf Skink		
Scincidae	<i>Morethia</i>	<i>lineocellata</i>	-		
Scincidae	<i>Morethia</i>	<i>obscura</i>	Woodland Flecked Skink		
Scincidae	<i>Tiliqua</i>	<i>rugosa rugosa</i>	Southwestern Bobtail		
Typhlopidae	<i>Ramphotyphlops</i>	<i>australis</i>	-		
Typhlopidae	<i>Ramphotyphlops</i>	<i>pinguis</i>	-		
Typhlopidae	<i>Ramphotyphlops</i>	<i>waitii</i>	-		
Varanidae	<i>Varanus</i>	<i>gouldii</i>	Gould's Sand Monitor		



Family	Genus	Species	Common Name	Status	Recorded During the Field Survey
Varanidae	<i>Varanus</i>	<i>tristis tristis</i>	Black-headed Monitor		
Amphibia					
Hylidae	<i>Litoria</i>	<i>adelaidensis</i>	Slender Tree Frog		
Hylidae	<i>Litoria</i>	<i>moorei</i>	Motorbike Frog or Bell Frog		
Myobatrachidae	<i>Crinia</i>	<i>georgiana</i>	Quacking Frog		
Myobatrachidae	<i>Crinia</i>	<i>glauerti</i>	Glauerts' Froglet		
Myobatrachidae	<i>Crinia</i>	<i>pseudinsignifera</i>	Bleating Froglet		
Myobatrachidae	<i>Geocrinia</i>	<i>leai</i>	Lea's Frog		
Myobatrachidae	<i>Heleioporus</i>	<i>albopunctatus</i>	Western Spotted Frog		
Myobatrachidae	<i>Heleioporus</i>	<i>barycragus</i>	Western Marsh Frog		
Myobatrachidae	<i>Heleioporus</i>	<i>eyrei</i>	Moaning Frog		
Myobatrachidae	<i>Heleioporus</i>	<i>inornatus</i>	Whooping Frog		
Myobatrachidae	<i>Heleioporus</i>	<i>psammophilus</i>	Sand Frog		
Myobatrachidae	<i>Limnodynastes</i>	<i>dorsalis</i>	Bullfrog or Banjo Frog		
Myobatrachidae	<i>Myobatrachus</i>	<i>gouldii</i>	Turtle Frog		
Myobatrachidae	<i>Pseudophryne</i>	<i>guentheri</i>	Crawling Frog or Gunther's Toadlet		
Fish					
Poeciliidae	<i>Gambusia</i>	<i>affinis</i>	Mosquito Fish / Plague Minnow	*	



GHD Pty Ltd ABN 39 008 488 373

GHD House, 239 Adelaide Tce. Perth, WA 6004



P.O. Box Y3106, Perth WA 6832

T: 61 8 6222 8222 F: 61 8 6222 8555 E: permail@ghd.com.au

© **GHD Pty Ltd 2007**

This document is and shall remain the property of GHD Pty Ltd. The document may only be used for the purposes for which it was commissioned and in accordance with the Terms of Engagement for the commission. Unauthorised use of this document in any form whatsoever is prohibited.

Document Status

Rev No.	Author	Reviewer		Approved for Issue		
		Name	Signature	Name	Signature	Date
	M Dilly	A Napier		M Goldstone		13/7/07